

Drugs that block allergy skin tests

*This is not an all inclusive list. Please call our office if you have any questions.

Allergy Eye Drops - Must be off 7 days prior: Such as Patanol, Pazeo, Pataday, Zaditor, Ketotifen generic.

Allergy Eye Drops - Must be off 4 weeks prior: Optivar

Nasal Sprays - Must be off 4 weeks prior: Astelin, Azelastine, Astepro, Patanase, Dymista.

Muscle Relaxants - Must be off 2 weeks prior: Flexeril, Cyclobenzaprine

Steroids - Prednisone 10mg or less.

Antihistamines - Must be off 7 days prior:

Actifed Cold and Allergy

Alavert

Allegra, Allegra-D 12 hour, Allegra-D 24 Hour

Astelin - **Must stop 4 weeks prior to testing**

Astepro - **Must stop 4 weeks prior to testing**

Atarax

Benadryl

Bromfed, Bromfed-DM, Bromfed-PD

brompheniramine

cetirizine

Clarinet, Clarinet Reditabs, Clarinet-D 12 Hour, Clarinet-D 24 Hour

Claritin, Claritin Reditabs, Claritin-D 12 Hour, Claritin-D 24 Hour Chlor-Trimeton

chlorpheniramine, chlorpheniramine/pseudoephedrine

clemastine fumarate

cyprohepatadine

Deconamine SR

Dimetapp, Dimetapp Cold & Allergy Elixir

diphenhydramine

Dramamine

fexofenadine

hydroxyzine

Ibuprofen Cold & Allergy

loratadine, loratadine/pseudoephedrine

Motrin Cold & Allergy

Palgic

Patanase - **Must stop for 4 weeks prior**

pseudoephedrine/triprolidine

Rondec DM Drops, Rondec DM Syrup, Rondec Drops, Rondec Syrup Semprex-D

Tavist Allergy

Tussionex PennKinetic

Tylenol Cold & Allergy

Vistaril

Xyzal

Zyrtec